

Effect of 12-week endurance training on biochemical parameters in elite football players: A comprehensive analysis

by Anonymous Anonymous

Submission date: 18-Oct-2023 06:56PM (UTC+0700)

Submission ID: 2183304065

File name: 10_OK.pdf (399.02K)

Word count: 4424

Character count: 25209

Effect of 12-week endurance training on biochemical parameters in elite football players: A comprehensive analysis

SM Farooque^{1abcde*}, Mukesh Mitra^{2abc}, & Prasanta Kumar Das^{1cde}

Tripura University (A Central University), India¹
MMDC College, India²

Received 24 July 2023; Accepted 26 September 2023; Published 19 October 2023
Ed 2023; 8(2): 388-395

ABSTRACT

Endurance training plays a pivotal role in football performance, shaping players' cardiovascular fitness and metabolic adjustments. The purpose of this study is to investigate the different changes in biochemical parameters after exposure to 12 weeks of circuit training based on endurance in football. A total of fifteen elite football players, with an average age of 16.87 ± 1.13 years, an average weight of 61.87 ± 5.94 kg and an average height of 172.82 ± 5.18 cm, participated in the study. On average, they had been training for 5.0 ± 1.0 years and had intermediate experience in national-level competitions for 4.0 ± 1.0 years. Following the initial data collection, the participants engaged in a twelve-week endurance training programme, involving one hour of daily training for four days a week. Pre-experimental designs were used in the data collection process. Further, for the analysis and extraction of data, descriptive statistics and t-tests were employed. The significance level was set at $P < 0.05$. Significant changes were found from baseline to post-treatment in each player's biochemical parameters. A significant increase in HDL, haemoglobin, creatine, and RBC and a significant decrease in LDL and VLDL were observed. The findings of this study underscore the potential benefits of endurance training for elite football players, as reflected in positive alterations in various biochemical parameters. However, for more comprehensive conclusions and a deeper understanding of the broader implications. Future research is recommended with larger sample sizes, control groups, and more detailed performance outcome measurements. This research contributes to our knowledge of the physiological responses of elite football players to endurance training and the potential advantages, it offers in optimising their performance and overall health.

Keywords: HDL; LDL; VLDL; creatine; hemoglobin



[https://doi.org/10.25299/sportarea.2023.vol8\(3\).13856](https://doi.org/10.25299/sportarea.2023.vol8(3).13856)

OPEN ACCESS



Copyright © 2023 SM Farooque, Mukesh Mitra, Prasanta Kumar Das

Corresponding Author: SM Farooque, ¹Department of Physical Education, Tripura University (A Central University), Agartala, India.

Email: smharish9@gmail.com

How to Cite: Farooque, SM., Mitra, M., & Das, P. K. (2023). Effect of 12-week endurance training on biochemical parameters in elite football players: A comprehensive analysis. *Journal Sport Area*, 8(3), 388-395. [https://doi.org/10.25299/sportarea.2023.vol8\(3\).13856](https://doi.org/10.25299/sportarea.2023.vol8(3).13856)

Authors' Contribution: a – Study Design; b – Data Collection; c – Statistical Analysis; d – Manuscript Preparation; e – Funds Collection

INTRODUCTION

Circuit training, a popular type of exercise, integrates strength training and cardiovascular exercise in a series of consecutive stations (Paoli et al., 2013; Mohammed et al., 2018). This dynamic approach to fitness

offers a range of benefits, including improved cardiovascular fitness, muscular strength and endurance (Fragala et al., 2019). However, while circuit training provides these benefits, its direct effects on certain blood components, such as red blood cells (RBCs), high-density lipoprotein (HDL), low-density lipoprotein (LDL), creatine, very low-density lipoprotein (VLDL), and haemoglobin, remain relatively limited (Kostrzewa-Nowak et al., 2020). Nevertheless, understanding these different effects may provide valuable insights into the holistic impact of circuit training on overall health and well-being.

Circuit training, a holistic exercise approach that combines strength and cardiovascular elements (Martin-Niedecken et al., 2020), may indirectly affect haemoglobin levels through improving cardiovascular fitness and oxygen transport capacity (Zivkovic et al., 2013). This is due to regular exercise can stimulate the production of more red blood cells and increase the efficiency of oxygen transport throughout the body, ultimately contributing to changes in performance patterns among athletes (Apostolidis et al., 2014). These interconnected physiological responses highlight the intricate relationship between circuit training, haemoglobin levels and athletic performance.

In the context of football, a sport that demands excellent fitness and endurance due to its dynamic nature, players undergo various forms of training to reach peak performance (Silva et al., 2023; Mujika et al., 2018). The rigorous demands go beyond daily training sessions, encompassing additional responsibilities such as national cups and other competitive events, creating a situation where football players experience significant levels of stress (McKay et al., 2022; Archiza et al., 2018). This co-occurring stress can put pressure on various physiological systems, including the musculoskeletal, neurological, immunological and metabolic systems, often resulting in observable changes in biochemical and haematological indices (Huggett, 2018). Therefore, the rigorous nature of the sport of football underscores the importance of understanding the interactions between training regimens, physiological responses and athletic performance.

Numerous epidemiological studies have consistently shown beneficial changes in blood lipid and lipoprotein levels resulting from regular aerobic exercise among both sexes (Sotiropoulos et al., 2009; Ruiz-Ramie et al., 2019; Martin-Niedecken et al., 2020). However, the lack of research dedicated to elite football athletes, particularly regarding their biochemical status, is noteworthy. Elite athletes engage in specialised training routines that may give rise to unique metabolic adaptations, potentially affecting their blood lipid profiles (Kostrzewa-Nowak et al., 2020). Thus, understanding specific lipid profile changes in elite players may provide insights to optimise training, nutrition and recovery strategies, which may ultimately improve cardiovascular health and longevity in the context of an intensive athletic career.

Furthermore, competitive stressors in sport can trigger physiological responses that affect the musculoskeletal, neurological, immunological and metabolic systems (Kostrzewa-Nowak et al., 2020). This can manifest as changes in stress-related biomarkers such as cortisol, serotonin, dopamine, creatine kinase (CK), lactate dehydrogenase (LDH), and myoglobin (Oglodek, 2022). Despite the recognition and admiration for football, limited research has explored the biochemistry and haematology of football players, especially regarding the physiological stressors imposed by the nature of the sport (Azmi & Kusnanik, 2018). Therefore, delving into this aspect has become very urgent to not only improve our knowledge of athletes' health, but also to develop better strategies in managing competitive stress that may significantly affect their performance.

Based on these considerations, this study investigated the impact of circuit training on biochemical parameters. Circuit training, which resembles high-intensity interval training, combines resistance and endurance training and involves a series of exercises performed at different stations (Silva et al., 2023). We conducted this study over half a competitive season on elite U-19 football players, aiming to deepen our understanding of their biochemical responses. The insights gained may prove valuable to coaches, physiotherapists and exercise experts, informing strategies to optimise training and performance among elite football players.

METHOD

The current study was carried out at Poloi Academy of Manipur, India, and involved fifteen elite football players who were part of the U-19 national team. These players had an average age of 16.87 ± 1.13 years, an

average weight of 61.87 ± 5.94 kg, and an average height of 172.82 ± 5.18 cm. They had a consistent track record of participating in competitions for the past 3-4 years and had also taken part in National Level Competitions. Before the training program commenced, the players were provided with detailed information about the study procedures, and informed consent was obtained from both the individuals and their parents through the academy. The study protocol was adhered to for each participant. Following the baseline data collection, all selected players underwent a 12-weeks endurance training program (Luo et al., 2023). The research took place between June and October 2022. The Institutional Human Ethical Committee approved the study protocol (Reference: TU/IHEC/3/1/22), ensuring compliance with the principles outlined in the declaration of Helsinki concerning human experimentation. The study focused on analyzing several biochemical variables, including red blood cells (RBC), creatine, hemoglobin, high-density lipoprotein (HDL), low-density lipoprotein (LDL), and very low-density lipoprotein (VLDL). The circuit training schedule program is represented in Fig 1.

Table 1. 12 Weeks Circuit Training Program on Elite Football Players

S. No	Exercise	Repetitions			Sets	Total duration
		1-4 weeks	5-8 weeks	9-12 weeks		
1.	Push up	25-30sec	35-40sec	40-45sec	3	1:00 hrs
2.	High knee	25-30sec	35-40sec	40-45sec	3	
3.	Plank	25-30sec	35-40sec	40-45sec	3	
4.	Burpee	25-30sec	35-40sec	40-45sec	3	
5.	Butt kick	25-30sec	35-40sec	40-45sec	3	
6.	Mountain climbing	25-30sec	35-40sec	40-45sec	3	
7.	Skipping	25-30sec	35-40sec	40-45sec	3	
8.	Sit up	25-30sec	35-40sec	40-45sec	3	




Figure 1. Schematic Diagram of Circuit Training Schedule

Procedure of Data Collection

For the blood collection, players were asked to place their forearm in a comfortable position. Before and after the training, a total of five milliliters of blood were drawn from the antecubital vein of each player

using two test tubes for the biochemical study, following the method described (Guo et al., 2022). Subsequently, the collected blood samples were used for the biochemical analysis.

Data Analysis

The data obtained from the study was analyzed using statistical software called SPSS version 27. Descriptive analysis, including calculations of mean, standard deviation (SD), and standard error mean (SEM) for each test, was conducted. Additionally, inferential statistics, specifically the pair sample t-test, were employed to determine the significance of differences between the Pre and Post tests.

Ethics

The study adhered to ethical guidelines outlined in the Declaration of Helsinki and received approval from the Institutional Human Ethical Committee (Reference: TU/IHEC/3/1/22). Prior to their involvement in the study, all participants or their legal guardians provided informed consent. Detailed information about the study protocol was provided to the participants, and they willingly signed the consent statement before participating in the study.

RESULTS AND DISCUSSION

The purpose of this study was to investigate how a 12-week circuit training program would influence both Biochemical Parameters and Skill Proficiency in U-19 elite football players. The researchers achieved this by comparing the baseline data with the data collected after the 12-week training period. The data collected from a group of 15 carefully selected subjects underwent comprehensive statistical analysis, and the findings of this study are elaborated upon below.

Table 2. Comparison on the Effect of Circuit Training on Biochemical Parameters of Football Player

Parameters	N	Test	Mean	SD	SEM	t-Value	p-Value
RBC	15	Pre test	5.57	.37	.09	3.51	.003
		Post test	5.65	.36	.09		
Creatine	15	Pre test	.91	.11	.02	3.59	.003
		Post test	.99	.10	.02		
Hemoglobin	15	Pre test	14.95	1.15	.29	3.09	.008
		Post test	15.27	1.19	.30		
HDL	15	Pre test	54.60	8.50	2.19	9.93	.000
		Post test	60.53	10.22	2.64		
LDL	15	Pre test	65.53	21.97	5.67	3.19	.006
		Post test	57.80	17.73	4.57		
VLDL	15	Pre test	16.20	4.57	1.49	6.34	.000
		Post test	13.73	4.16	1.32		

The effect of circuit training on RBC evoked a significant increase from pre to post data ($5.57 \pm .37$) to ($5.65 \pm .36$) which shown a t value of 3.51 having p-Value of .003 which is significant at .003. When we compare with this pre value with biological reference frame by the WHO (4.5-5.5) reference value, the effect in the changes has surpass the borderline of RBC which need a concern on this specific variable for further studies. The same changes have also observed on creatine variables after 12 weeks circuit training.

The mean and SD observed in the pre were ($.91 \pm .11$) which further increase to ($.99 \pm .10$) after long 12 weeks circuit training. In addition, significant changes were observed when we compared the pre data to post data. The observed t value in the comparison was 3.59 which is significant at ($p = .003$). The hemoglobin changes depicted from the training of circuit training of 12 weeks was (14.95 ± 1.15) pre and (15.27 ± 1.19) post. And the significant difference found between pre and post were $p = .008$ having a t value of 3.09 which is significant at 0.05 significant level.

Evidence shown by the table 4.2 on HDL shown a significant increase from pre to post (54.60 ± 8.50) and (60.53 ± 10.22). Further shown a significant t value of 9.93 in compare to pre and post which is significant at 0.05 level. The study shown a significant reduction in the mean value from pre to post in case of LDL which

shown positive effect on the changes of biochemical variables. The mean value of LDL from pre to post were (65.53 ± 21.97) and (57.80 ± 17.73) respectively. The result has been compared to find the significant difference and the t value of 3.19 was observed with significant value of $p = .006$. The mean value and Sd observed for pre and post of VLDL were (16.20 ± 4.57) and (13.73 ± 4.16) respectively. In addition, significant difference was also observed at 0.05 level of significance.

The findings of this study suggest that the circuit training programme induced significant changes in selected variables, including various physiological alterations that led to improved individual performance. Red blood cells (RBCs) play a crucial role in microcirculation by facilitating the exchange of oxygen and carbon dioxide with surrounding tissues (Beunders et al., 2020; Gates et al., 2013). To navigate microscopic capillaries, RBCs must deform, and their primary function is to transport respiratory gases. During exercise, the circulatory system ensures the supply of substrates to working muscles. The primary role of RBCs during exercise is to transport oxygen from the lungs to tissues in response to the body's demands and to transport metabolically produced carbon dioxide to the lungs for elimination (Mairbäurl, 2013; Kostorzewa-Nowak et al., 2015). They also contribute to energy production and repair. This study observed significant changes in red blood cell counts in both training methods. Untrained healthy adults, as per Calahorra (2011), can benefit from a regular moderate running training programme due to the regeneration of a younger RBC population with improved characteristics, aiding oxygen supply in the microcirculation.

Creatinine levels in the human body vary among individuals. Individuals engaged in moderate or intense physical exercise typically exhibit higher serum creatinine levels than sedentary individuals (Zivkovic et al., 2013). Factors such as age, gender, ethnicity, dietary protein intake, lean mass, and intensity of physical activity can influence creatinine concentrations, either decreasing or increasing them (Arwandi & Firdaus, 2021; Beunders et al., 2020). The current study noted a substantial increase in creatine levels in football players after a 12-week circuit training program. This increase can be attributed to the various stress-induced muscle changes that result in muscle tear and injury, with creatinine being expelled as a waste product through the kidneys (Bompa & Buzzichelli, 2019). Muscle requires a certain amount of creatine for energy synthesis to adapt to training-induced stress and increased loads.

The study's alignment with Ruiz-Ramie et al. (2019) findings underscores the consistency of the impact of light to moderate exercise on net protein catabolism and creatinine excretion. Additionally, this perspective on creatine concentration, as introduced in the presented study, sheds light on the significance of monitoring creatinine expulsion as an indicator of metabolic adaptations in response to exercise (Asadi, 2016). These insights contribute to our understanding of how exercise-induced stress influences metabolic processes and further emphasise the relevance of creatinine as a biomarker for muscular stress and adaptation (Guo et al., 2022).

Exercise has the potential to elevate haemoglobin levels and improve the overall quality of life. Although this study focused on women, it addressed a specific community of male U-19 high-level players. The observation of RBCs suggests a likely favourable effect on haemoglobin concentration due to their direct association. As (Törpel et al., 2020) suggests, an increase in RBC volume leads to a higher haemoglobin concentration in the blood, enhancing oxygenated blood distribution in the body and supporting various bodily functions. The current study indicates a trend of rising RBC levels after 12 weeks of circuit training for football players, suggesting a concurrent increase in haemoglobin levels.

Athletes serve as models of HDL regulation due to their elevated HDL levels (Manna et al., 2010). Low HDL concentrations have been identified as a risk factor for cardiac diseases, with a 1 mg/dl increase in HDL lowering heart disease risk by 4% (Gordon et al., 2014). Notably, high-profile players may experience unexpected attacks during matches for unknown reasons. According to Osei et al. (2020), short-term, low-intensity exercise can influence HDL concentration, but unaltered evidence also exists. It has been reported that prolonged activity exceeding 1.5 hours is required for HDL changes to occur during, immediately after, and the day following exercise. In contrast, the present study's statistical data indicate a significant increase in HDL levels in football players after a 12-week circuit training programme, highlighting a pronounced shift in the pattern of HDL levels among these athletes (Sepriadi et al., 2020).

CONCLUSION

The 12-week endurance training programme has shown promise as an avenue for enhancing the performance potential of elite football players. The observed favourable changes in various biochemical markers suggest that focused endurance training can significantly contribute to improving the overall athletic capacity of these players. However, this study acknowledges certain limitations, including the relatively small sample size and the absence of a control group, which may affect the generalizability and causality of the findings. Moreover, individual variations in lifestyle and dietary habits among participants introduce confounding factors that warrant further investigation. Therefore, it is recommended that future research endeavors involve larger-scale studies with diverse participant pools, controlled trials to establish causality, and longitudinal studies to assess the sustainability of the observed changes. This research contributes to our understanding of the potential benefits of endurance-based circuit training for elite football players and underscores the need for further clinical research to precisely elucidate its role in enhancing individual performance and skill development within the professional sports context.

ACKNOWLEDGEMENTS

The author of the study is deeply grateful to ¹ Dr. Krishendu Dhar, Assistant Professor in the Department of Physical Education at Tripura University, who has been a tremendous source of knowledge and inspiration. The author holds great appreciation for Dr. Sudip Das, Associate Professor in the same department, for providing invaluable advice despite his busy schedule. The technical assistance provided by Dr. Sanjib Kumar Bhowmik, Assistant Professor in the department of physical education at Tripura University, is also acknowledged with gratitude. Additionally, the authors express their appreciation to the previous author for offering them the opportunity to expand their understanding of the subject matter.

CONFLICT OF INTEREST

¹ The authors declare no conflict of interest.

REFERENCES

- Apostolidis, N., Bogdanis, G. C., Kostopoulos, N., Souglis, A., & Papadopoulos, Ch. (2014). Changes in the Lipid Profile of Elite Basketball and Soccer Players after a Match. *Research in Sports Medicine*, 22(1), 100–110. <https://doi.org/10.1080/15438627.2013.852095>
- Archiza, B., Andaku, D. K., Caruso, F. C. R., Bonjorno, J. C., Oliveira, C. R. de, Ricci, P. A., Amaral, A. C. do, Mattiello, S. M., Libardi, C. A., Phillips, S. A., Arena, R., & Borghi-Silva, A. (2018). Effects of Inspiratory Muscle Training in Professional Women Football Players: A Randomized Sham-Controlled Trial. *Journal of Sports Sciences*, 36(7), 771–780. <https://doi.org/10.1080/02640414.2017.1340659>
- Arwandi, J., & Firdaus, M. (2021). Effect of Agility Training Towards Soccer Dribbling Skills. *Proceedings of the 1st International Conference on Sport Sciences, Health and Tourism (ICSSHT 2019)*. <https://doi.org/10.2991/ahsr.k.210130.002>
- Asadi, A. (2016). Relationship Between Jumping Ability, Agility and Sprint Performance of Elite Young Basketball Players: A Field-Test Approach. *Brazilian Journal of Kinanthropometry and Human Performance*, 18(2), 177-186. <https://doi.org/10.5007/1980-0037.2016v18n2p177>
- Azmi, K., & Kusnanik, N. W. (2018). Effect of Exercise Program Speed, Agility, and Quickness (SAQ) in Improving Speed, Agility, and Acceleration. *Journal of Physics: Conference Series*, 947, 012043. <https://doi.org/10.1088/1742-6596/947/1/012043>
- Beunders, R., Bongers, C. C. W. G., & Pickkers, P. (2020). The Effects of Physical Exercise on the Assessment of Kidney Function. *Journal of Applied Physiology*, 128(5), 1459–1460. <https://doi.org/10.1152/jappphysiol.00189.2020>

- Bompa, T. O. , & Buzzichelli. C. (2019). *Periodization Theory and Methodology of Training* (Sixth). Human Kinetics
- Fragala, M. S., Cadore, E. L., Dorgo, S., Izquierdo, M., Kraemer, W. J., Peterson, M. D., & Ryan, E. D. (2019). Resistance Training for Older Adults: Position Statement From the National Strength and Conditioning Association. *Journal of Strength and Conditioning Research*, 33(8), 2019–2052. <https://doi.org/10.1519/JSC.0000000000003230>
- Gates, N., Fiatarone Singh, M. A., Sachdev, P. S., & Valenzuela, M. (2013). The Effect of Exercise Training on Cognitive Function in Older Adults with Mild Cognitive Impairment: A Meta-analysis of Randomized Controlled Trials. *The American Journal of Geriatric Psychiatry*, 21(11), 1086–1097. <https://doi.org/10.1016/j.jagp.2013.02.018>
- Gordon, B., Chen, S., & Durstine, J. L. (2014). The Effects of Exercise Training on the Traditional Lipid Profile and Beyond. *Current Sports Medicine Reports*, 13(4), 253–259. <https://doi.org/10.1249/JSR.0000000000000073>
- Guo, Z., Cai, J., Wu, Z., & Gong, W. (2022). Effect of High-Intensity Interval Training Combined with Fasting in the Treatment of Overweight and Obese Adults: A Systematic Review and Meta-Analysis. *International Journal of Environmental Research and Public Health*, 19(8), 4638. <https://doi.org/10.3390/ijerph19084638>
- Huggett, R. J. (2018). *Biomarkers: Biochemical, Physiological, and Histological Markers of Anthropogenic Stress* (CRC Press, Ed.).
- Kostrzewa-Nowak, D., Nowak, R., Jastrzębski, Z., Zarębska, A., Bichowska, M., Drobnik-Kozakiewicz, I., Radziwiński, Ł., Leońska-Duniec, A., Ficek, K., & Cięszczyk, P. (2015). Effect of 12-Week-Long Aerobic Training Programme on Body Composition, Aerobic Capacity, Complete Blood Count and Blood Lipid Profile among Young Women. *Biochemia Medica*, 103–113. <https://doi.org/10.11613/BM.2015.013>
- Kostrzewa-Nowak, D., Nowakowska, A., Zwierko, T., Rybak, M., & Nowak, R. (2020). The Influence of a Health-Related Fitness Training Program on Motor Performance as Well as Hematological and Biochemical Parameters. *International Journal of Environmental Research and Public Health*, 17(2), 578. <https://doi.org/10.3390/ijerph17020578>
- Luo, S., Soh, K. G., Zhang, L., Zhai, X., Sunardi, J., Gao, Y., & Sun, H. (2023). Effect of Core Training on Skill-Related Physical Fitness Performance among Soccer Players: A Systematic Review. *Frontiers in Public Health*, 10. <https://doi.org/10.3389/fpubh.2022.1046456>
- Mairbäurl, H. (2013). Red Blood Cells in Sports: Effects of Exercise and Training on Oxygen Supply by Red Blood Cells. *Frontiers in Physiology*, 4. <https://doi.org/10.3389/fphys.2013.00332>
- Manna, I., Khanna, G. L., & Dhara, P. C. (2010). Variation in Age and Training on Selected Biochemical Variables of Indian Hockey Players. *Al Ameen J Med Sci*, 3(2), 128-139.
- Martin-Niedecken, A. L., Mahrer, A., Rogers, K., de Bruin, E. D., & Schättin, A. (2020). “HIIT” the ExerCube: Comparing the Effectiveness of Functional High-Intensity Interval Training in Conventional vs. Exergame-Based Training. *Frontiers in Computer Science*, 2. <https://doi.org/10.3389/fcomp.2020.00033>
- McKay, A., Cropley, B., Mullen, R., Shearer, D., & Hanton, S. (2022). Psychosocial Demands and Situational Properties of the Club-to-International Transition in Male Youth Football. *Journal of Applied Sport Psychology*, 34(6), 1272–1294. <https://doi.org/10.1080/10413200.2021.1972495>

- Mohammed, Z., Zohar, B. F., Gourar, B., Ali, B., & Idriss, M. M. (2018). VO₂max Levels as a Pointer of Physiological Training Status among Soccer Players. *Acta Facultatis Educationis Physicae Universitatis Comenianae*, 58(2), 112–121. <https://doi.org/10.2478/afepuc-2018-0010>
- Mujika, I., Halson, S., Burke, L. M., Balagué, G., & Farrow, D. (2018). An Integrated, Multifactorial Approach to Periodization for Optimal Performance in Individual and Team Sports. *International Journal of Sports Physiology and Performance*, 13(5), 538–561. <https://doi.org/10.1123/ijssp.2018-0093>
- Ogłodek, E. A. (2022). Changes in the Serum Concentration Levels of Serotonin, Tryptophan and Cortisol among Stress-Resilient and Stress-Susceptible Individuals after Experiencing Traumatic Stress. *International Journal of Environmental Research and Public Health*, 19(24), 16517. <https://doi.org/10.3390/ijerph192416517>
- Osei, F., Moses, M. O., Pambo, P., Baffour-Awuah, B., Asamoah, B., Afrifa, D., Appiah, E. J., Akwa, L. G., & Obour, A. (2020). Changes in Cardiovascular Parameters of a-University Football athletes Associated with Short duration Pre-Tournament Training. *Scientific African*, 8, e00285. <https://doi.org/10.1016/j.sciaf.2020.e00285>
- Paoli, A., Pacelli, Q. F., Moro, T., Marcolin, G., Neri, M., Battaglia, G., Sergi, G., Bolzetta, F., & Bianco, A. (2013). Effects of High-Intensity Circuit Training, Low-Intensity Circuit Training and Endurance Training on Blood Pressure and Lipoproteins in Middle-Aged Overweight Men. *Lipids in Health and Disease*, 12(1). <https://doi.org/10.1186/1476-511X-12-131>
- Ruiz-Ramie, J. J., Barber, J. L., & Sarzynski, M. A. (2019). Effects of Exercise on HDL Functionality. *Current Opinion in Lipidology*, 30(1), 16–23. <https://doi.org/10.1097/MOL.0000000000000568>
- Sepriadi, Jannah, K., & Eldawaty. (2020). The Effect of Jogging Exercise to Improve Hemoglobin Levels. *Journal of Physics: Conference Series*, 1481(1), 012028. <https://doi.org/10.1088/1742-6596/1481/1/012028>
- Silva, H., Nakamura, F. Y., Beato, M., & Marcelino, R. (2023). Acceleration and Deceleration Demands during Training Sessions in Football: A Systematic Review. *Science and Medicine in Football*, 7(3), 198–213. <https://doi.org/10.1080/24733938.2022.2090600>
- Sotiropoulos, A., Travlos, A. K., Gissis, I., Souglis, A. G., & Grezios, A. (2009). The Effect of a 4-Week Training Regimen on Body Fat and Aerobic Capacity of Professional Soccer Players During The Transition Period. *Journal of Strength and Conditioning Research*, 23(6), 1697–1703. <https://doi.org/10.1519/JSC.0b013e3181b3df69>
- Törpel, A., Peter, B., & Schega, L. (2020). Effect of Resistance Training Under Normobaric Hypoxia on Physical Performance, Hematological Parameters, and Body Composition in Young and Older People. *Frontiers in Physiology*, 11. <https://doi.org/10.3389/fphys.2020.00335>
- Zivkovic, V., Lazarevic, P., Djuric, D., Cubrilo, D., Macura, M., Vuletic, M., Barudic, N., Nestic, M., & Jakovljevic, V. (2013). Alteration in Basal Redox State of Young Male Soccer Players after a Six-Month Training Programme. *Acta Physiologica Hungarica*, 100(1), 64–76. <https://doi.org/10.1556/APhysiol.100.2013.1.6>

Effect of 12-week endurance training on biochemical parameters in elite football players: A comprehensive analysis

ORIGINALITY REPORT

9%

SIMILARITY INDEX

0%

INTERNET SOURCES

0%

PUBLICATIONS

9%

STUDENT PAPERS

PRIMARY SOURCES

1

Submitted to Eastern Arizona College

Student Paper

9%

Exclude quotes Off

Exclude matches < 5%

Exclude bibliography On