

The relationship between subjective knee function and psychological readiness to return to sport after ACL injury: A cross-sectional study

Ela Yuliana^{1acde,*} , Ruliando Hasea Purba^{1abc} , Bazzar Ari Migrha^{1abc} ,
Nadya Dwi Oktafiranda^{1abc} , & Dewi Nurhidayah^{2abde} 

Universitas Negeri Jakarta, Indonesia¹
Universitas Cenderawasih, Indonesia²

Received 25 October 2025; Accepted 15 November 2025; Published 14 December 2025
Ed 2025; 10(3): 369-376

ABSTRACT

Background: Athletes with anterior cruciate ligament (ACL) injuries often reduce or stop sports participation due to fear of re-injury and low confidence in their injured knee. While most studies emphasize physical recovery, fewer explore how subjective knee function affects psychological readiness to return to sport. **Objectives:** This study aims to determine the contribution of self-reported knee symptoms and function (subjective assessment) to psychological readiness to return to sport after ACL injury with operative and non-operative treatment. **Methods:** This cross-sectional study involved 40 participants who met the inclusion criteria for post-ACL injury, both operative and non-operative. Data were collected using the International Knee Documentation Committee (IKDC 2000) questionnaire for subjective knee function and the Anterior Cruciate Ligament–Return to Sport after Injury (ACL-RSI) scale for psychological readiness. A simple linear regression test was conducted with a 95% confidence interval. **Finding/Results:** The mean IKDC score was 69.80 ± 12.53 and the mean ACL-RSI score was 58.43 ± 19.12 . The regression analysis showed a significant positive relationship between subjective knee function and psychological readiness to return to sport ($r = 0.61$, $p < 0.05$), with a coefficient of determination (R^2) of 0.373. **Conclusion:** Subjective knee function significantly influences psychological readiness to return to sport after an ACL injury. Enhancing knee function through targeted rehabilitation programs may improve athletes' confidence and reduce fear of re-injury, facilitating a safer and more effective return to sport.

Keywords: Anterior cruciate ligament injury; knee function; psychological readiness

 [https://doi.org/10.25299/sportarea.2025.vol10\(3\).19470](https://doi.org/10.25299/sportarea.2025.vol10(3).19470)

OPEN ACCESS



Copyright © 2025 Ela Yuliana, Ruliando Hasea Purba, Bazzar Ari Migrha, Nadya Dwi Oktafiranda, Dewi Nurhidayah

 **Corresponding Author:** Ela Yuliana, Department of Sports Science, Faculty of Sports Science, Universitas Negeri Jakarta, Jakarta, Indonesia
 ela_yuliana@unj.ac.id

How to Cite: Yuliana, E., Purba, R. H., Migrha, B. A., Oktafiranda, N. D., & Nurhidayah, D. (2025). The relationship between subjective knee function and psychological readiness to return to sport after ACL injury: A cross-sectional study. *Journal Sport Area*, 10(3), 369-376. [https://doi.org/10.25299/sportarea.2025.vol10\(3\).19470](https://doi.org/10.25299/sportarea.2025.vol10(3).19470)

Authors' Contribution: a – Study Design; b – Data Collection; c – Statistical Analysis; d – Manuscript Preparation; e – Funds Collection

INTRODUCTION

ACL injuries type of musculoskeletal injury that can have significant impacts, especially for adolescents who are actively involved in competitive sports. This injury not only limits a person's ability to return to previous performance but can also potentially stop sports participation completely. Previous research reported

that only two out of five athletes successfully returned to the same sport as before the injury two years after ACL reconstruction surgery, or approximately 60% (Ardern et al., 2015). Similar results were reported by Lindanger et al. (2019) as many as 83% of patients returned to pivot sports, while 53% managed to return to their previous level of performance after ACL reconstruction surgery. Supported by a recent study comparing physical activity levels between uninjured individuals and individuals after ACL reconstruction, individuals observed 5-8 years post ACL reconstruction surgery have a physical activity is lower than those who have never been injured (Stigert et al., 2023). This condition can have long-term impacts, namely an increased risk of osteoarthritis (OA) and if individuals already have OA, individuals with a history of ACL injury have a seven-fold increased risk of undergoing total knee replacement (TKR) (Khan et al., 2019).

Research suggests that psychological components are associated with individuals' restrictions on returning to exercise after an ACL injury, as various emotional and cognitive reactions can arise after an ACL injury, like afraid of getting injured again (76.7%), deficiency of confidence in the abilities of the injured knee (14.8%), depression (5.6%), and decreased interest or motivation after the injury (2.5%) (Nwachukwu et al., 2019). Meanwhile, psychological readiness is closely related and significantly influences an individual's return to physical activity as previous the ACL injury, so that the psychological readiness score is a predictor of success or failure in returning to exercise after an ACL injury (Beischer et al., 2023; Vutescu et al., 2021; Webster & Feller, 2019). Patients who successfully return to the same physical activities after ACL reconstruction surgery are generally characterized by a higher level of psychological readiness (Beischer et al., 2023). Supported by other studies, higher psychological readiness scores are associated with a return to pre-injury performance (Vutescu et al., 2021; Webster & Feller, 2019). Conversely, those with lower physical activity levels compared to before the ACL injury showed lower psychological readiness scores (Yuliana & Kushartanti, 2020). Therefore, low psychological readiness scores are an indicator of failure to return to post ACL sport (Faleide et al., 2021; Schilaty et al., 2023).

Many factors can influence psychological readiness to return to sport after an ACL injury, including physical condition (Faleide et al., 2021), gender, age, time between injury and surgery, pre-injury sports participation, and knee function (Webster et al., 2018). Research findings indicate that subjective knee symptoms and function are closely related to psychological readiness to return to sport after ACL surgery. (Webster et al., 2018). Subjective knee symptoms and function are self-reported assessments by patients through administered questionnaires. Strengthened by a biopsychosocial approach, self-reported knee function and psychological readiness to return to sport are biopsychosocial variables that influence return to sport after sports injury (Bae, 2024). The biopsychosocial approach is how biological, psychological and social factors influence the occurrence of sports injuries and the results of sports injury rehabilitation (Bae, 2024). However, limited research in Indonesia has examined how subjective knee function relates to psychological readiness after ACL injury. Therefore, this study aims to examine the relationship between these two factors among post-ACL athletes.

METHOD

Research Design

This study used a cross-sectional survey design, and has been declared to meet the requirements or be ethically feasible with the number KEP/II/2024/X/M290524EY-PMNT.

Research Subject

This study sample involved 40 individuals with ACL injuries treated either operatively or non-operatively. The sample met the criteria established in this study. Inclusion criteria included experiencing an ACL injury due to sports activities, and a time interval of more than six months from the ACL injury or ACL reconstruction surgery. This time period is a common measure used to assess knee function after ACL surgery (Kay et al., 2023; Nawasreh et al., 2018). Furthermore, participants were willing to participate in the study by signing a consent agreement to be included in the study. Meanwhile, exclusion criteria aimed to avoid influencing the reported knee function assessment results and causing bias in the relationship to psychological readiness, including experiencing other injuries besides the ACL injury after or before the ACL injury, ongoing signs of

inflammation in the knee joint, and range of motion still below 50% compared to the normal side. Furthermore, participants had health problems or conditions that interfered with giving consent to be included in the study, such as problems with the neuromuscular system or psychological problems, thus not representing the target population. Of course, the determination of research sample criteria is adjusted based on the research objectives so that the selected sample will be able to provide relevant and accurate information to answer the research questions.

Research Procedure

This study was conducted in July 2024, at the Faculty of Sport Science, Jakarta State University. Subjective knee symptoms and function were assessed using the IKDC 2000 subjective knee joint assessment form, and psychological readiness was assessed using the ACL-RSI questionnaire, translated into Indonesian using the Cross-Cultural Adaptation method (Yuliana & Kushartanti, 2020). The IKDC 2000 instrument consists of 18 questions, scored between 0 and 100, with higher scores indicating lower levels of symptoms and functional impairment and higher levels of sports activity (Kanakamedala et al., 2016). Meanwhile, the ACL-RSI questionnaire consists of 12 questions and was developed based on three components related to returning to sport (emotion, performance confidence, and risk assessment) and scored between 0 and 100, with higher scores indicating a more positive psychological state (Sadeqi et al., 2018). The research sample was asked to answer the IKDC 2000 questionnaire according to the symptoms and knee function currently felt, and to answer the ACL-RSI questionnaire according to their current psychological condition.

Statistic Analysis

This research analyzes data using Statistical Package for the Social Sciences (SPSS) Version 20 software. Descriptive statistics were used to analyze the baseline characteristics of the study sample. Data included the mean, maximum, minimum, and standard deviation, with 95% confidence intervals (CI). Data normality was tested using the Shapiro–Wilk test. A linearity test was performed to test the linear regression assumption. Furthermore, a simple linear regression test was used to determine the correlation coefficient and coefficient of determination of subjective knee symptoms and function, as well as psychological readiness, with a significance level of 0.05. Interpretation criteria for correlation strength $r > 0.6$ = strong.

RESULTS AND DISCUSSION

A total of 40 research samples participated and met the criteria set in this study. Consisting of 31 male samples (77.5%) and 9 female samples (22.5%). The research sample with operative treatment consisted of 26 samples (65%) and non-operative treatment consisted of 14 samples (35%). Characteristics data of the research sample are in **Table 1**.

Table 1. Research Subject Characteristics Data

No	Characteristics	Mean	Standard Deviation
1	Age	22.4	2.76
2	Time Between Surgery and Test (operative treatment)*	31.25	20.52
3	Time Between Injury and Test (non-operative treatment)*	28.77	18.06

Note: *units using months

The research sample of 40 with operative or non-operative treatment had an average IKDC score was 69.80 ± 12.53 (95% CI 65.79 to 73.81), and the average ACL-RSI was 58.43 ± 19.12 (95% CI 52.31 to 64.54). The assessment results are presented in **Table 2**.

Table 2. IKDC 2000 Score and ACL-RSI Score Results

No	Description	Mean	Maximum	Minimum	Standard Deviation	95% Confidence Interval of the Difference	
						Lower	Upper
1	IKDC 2000	69.80	95.40	19.5	12.53	65.79	73.82
2	ACL-RSI	58.43	97	21	19.12	52.31	64.54

A significant positive correlation was found between IKDC scores and ACL-RSI ($r = 0.611$, $p < 0.05$). Furthermore, to determine the extent of influence of subjective knee symptoms and function on psychological readiness to return to exercise post an ACL injury, a coefficient of determination test was conducted. The results showed an R Square value (coefficient of determination) of 0.373. This means that subjective knee symptoms and function have an influence or contribution to psychological readiness to return to exercise post an ACL injury of 37.3%. In addition, a correlation coefficient analysis was also conducted between scores IKDC 2000 and scores ACL-RSI. The results showed a figure of 0.61, which means that subjective knee symptoms and function have a positive and strong relationship with psychological readiness to return to exercise after an ACL injury. The results of the data analysis are presented in **Table 3**.

Table 3. Simple Linear Regression Test Results

Independent Variables	R (Correlation Coefficient)	R Square (Determination Coefficient)	P-Value
IKDC 2000 Score (Subjective Knee Symptoms and Function)	0.611	0.373	0.000

Psychological readiness, although not the primary focus in ACL injury rehabilitation, has a significant impact on recovery outcomes after an ACL injury (Nedder et al., 2025). Nearly two-thirds (60% of athletes) are not psychologically prepared to return to sport after injury, and only 40% are ready to return to sport after an ACL injury (Juggath & Naidoo, 2024). Psychological readiness is defined as feelings of confidence and the absence of fear (Dluzniewski et al., 2024). The purpose of this study was to examine the relationship between subjective knee function with psychological readiness to return to sport after an ACL injury.

The Relationship between Knee Function and Psychology

A significant relationship was found between subjective knee symptoms and function as assessed by the IKDC 2000 and psychological readiness to return to sport after an ACL injury as evaluated by the ACL-RSI. Consistent with previous research, Psychological readiness that is highly correlated with returning to sport after ACL surgery is self-reported symptoms and function (Schilaty et al., 2023; Webster et al., 2018). Furthermore, subjective knee symptoms and function contributed 37.3% to psychological readiness to return to sport after an ACL injury. This supports previous research findings, where quadriceps and hamstring rate of torque development (RTD) have been identified as determinants of psychological readiness to return to sport after an ACL injury (Lee et al., 2023). Conversely, patients with lower functional assessment results, such as hamstring strength, jumping performance, and subjective knee function, had higher kinesiophobia (Norte et al., 2019).

The Role of Muscle Strength

To explain the relationship between knee symptoms and function and psychological readiness, as demonstrated in the results of this research and supported by several previous studies, when an ACL injury or post-ACL reconstruction occurs, common problems found are atrophy and deficits in quadriceps muscle strength and endurance (Csapo et al., 2019; Pamukoff et al., 2017). Furthermore, ACL injury causes deficits in hamstring muscle strength (Bouzekraoui Alaoui et al., 2025). Hamstring muscle strength that fails to recover can increase the risk of re-injury, possibly due to the inability to support and stabilize the knee joint during changes in frontal and transverse planes of movement (Buckthorpe et al., 2021). Deficits in hamstring and quadriceps muscle strength can also lead to the perception of joint instability during sports participation (Lee et al., 2023), and movement becomes impaired, thus affecting the athlete’s movement quality (Buckthorpe, 2019). This is likely to be perceived negatively by athletes, thus altering ACL-RSI scores (Lee et al., 2023). This is in line with the statement that functional decline, such as knee instability after an ACL injury, can trigger negative psychological responses (Christino et al., 2016). When negative psychological responses arise, they can become a barrier to successful motor and functional recovery after ACL surgery (Du et al., 2022). Therefore, psychological readiness scores can be used as predictors of a person’s safe return to sport post injury (Dluzniewski et al., 2024).

Implications for Rehabilitation

The return of knee function to its original state and the return of physical activity or sports as before the injury are signs of successful ACL injury treatment, both with operative and non-operative treatment. Therefore, understanding interventions or rehabilitation programs that can improve knee function is necessary to minimize the fear of ACL re-injury and increase psychological readiness to return to sport after an ACL injury. Rehabilitation programs have been shown to not only improve function but also improve athlete psychology (Meierbachtol et al., 2018). A recent study reported that patients who underwent symmetrical muscle function restoration 12 months after ACL reconstruction had higher self-efficacy compared to patients who did not undergo knee function recovery (Nedder et al., 2025). Athletes who participated in advanced recovery training after ACL reconstruction surgery not only experienced improved knee function but also improved athlete psychology (Meierbachtol et al., 2018). Similarly, a more stable knee can provide a feeling of greater confidence and support better recovery outcomes (Christino et al., 2016). These data support the relationship between physical ability or performance and cognitive and emotional responses, such as be afraid and confidence, then contribute to athletes' perceptions of skill or success in sport participation (Arden et al., 2013). Therefore, in ACL injury rehabilitation interventions, physical and psychological recovery are inseparable and are part of the achievement indicators, in order to optimize the results of the ACL injury rehabilitation program and increase the chances of a safe return to sport. Furthermore, knee symptoms and function, as well as psychological readiness, are crucial factors that can play a part in athletes' decisions to return to sport.

Our study has limitations, including the fact that the data analysis did not differentiate between operative and non-operative treatment groups. Future research could analyze these groups by operative and non-operative treatment, which would provide broader context for the findings. Another limitation is that this study only highlights the relationship between self-reported knee symptoms and function and psychological readiness to return to sport after ACL injury. Future research may identify factors such as quadriceps-hamstrings strength, biomechanics, neuromuscular control, and joint stability on psychological readiness to return to sport post-ACL injury.

CONCLUSION

Subjective knee function significantly influences psychological readiness to return to sport following ACL injury. Rehabilitation programs that enhance knee function may consequently improve athletes' confidence and reduce fear of re-injury.

ACKNOWLEDGEMENTS

The researchers would like to thank the student athletes from the Faculty of Sport Science, Jakarta State University, who agreed to be research subjects. They also thank the Faculty of Sport Science, Jakarta State University, for their contribution to this research, which provided a 2024 research grant to the researchers, making this research possible.

REFERENCES

- Arden, C. L., Taylor, N. F., Feller, J. A., & Webster, K. E. (2013). A Systematic Review of the Psychological Factors Associated with Returning to Sport Following Injury. *British Journal of Sports Medicine*, 47(17), 1120–1126. <https://doi.org/10.1136/bjsports-2012-091203>
- Arden, C. L., Taylor, N. F., Feller, J. A., Whitehead, T. S., & Webster, K. E. (2015). Sports Participation 2 Years after Anterior Cruciate Ligament Reconstruction in Athletes who Had not Returned to Sport At 1 Year: A Prospective Follow-Up of Physical Function and Psychological Factors in 122 Athletes. *American Journal of Sports Medicine*, 43(4), 848–856. <https://doi.org/10.1177/0363546514563282>
- Bae, M. (2024). Biopsychosocial Approach to Sports Injury: A Systematic Review and Exploration of Knowledge Structure. *BMC Sports Science, Medicine and Rehabilitation*, 16(1), 1-11. <https://doi.org/10.1186/s13102-024-01025-x>

- Beischer, S., Hamrin Senorski, E., & Thomeé, R. (2023). Patients That Maintain Their Pre-Injury Level of Physical Activity 3–5 Years after ACL Reconstruction are, 18 Months after Surgery, Characterised by Higher Levels of Readiness to Return to Sport. *Knee Surgery, Sports Traumatology, Arthroscopy*, 31(2), 596–607. <https://doi.org/10.1007/s00167-022-07230-w>
- Bouzekraoui Alaoui, I., Moiroux-Sahraoui, A., Mazeas, J., Kakavas, G., Bialy, M., Douryang, M., & Forelli, F. (2025). Impact of Hamstring Graft on Hamstring Peak Torque and Maximum Effective Angle After Anterior Cruciate Ligament Reconstruction: An Exploratory and Preliminary Study. *Bioengineering*, 12(5), 465. <https://doi.org/10.3390/bioengineering12050465>
- Buckthorpe, M. (2019). Optimising the Late-Stage Rehabilitation and Return-to-Sport Training and Testing Process After ACL Reconstruction. *Sports medicine (Auckland, N.Z.)*, 49(7), 1043–1058. <https://doi.org/10.1007/s40279-019-01102-z>
- Buckthorpe, M., Danelon, F., La Rosa, G., Nanni, G., Stride, M., & Della Villa, F. (2021). Recommendations for Hamstring Function Recovery after ACL Reconstruction. *Sports medicine (Auckland, N.Z.)*, 51(4), 607–624. <https://doi.org/10.1007/s40279-020-01400-x>
- Christino, M. A., Fleming, B. C., Machan, J. T., & Shalvoy, R. M. (2016). Psychological Factors Associated With Anterior Cruciate Ligament Reconstruction Recovery. *Orthopaedic Journal of Sports Medicine*, 4(3), 1-9. <https://doi.org/10.1177/2325967116638341>
- Csapo, R., Hoser, C., Gföller, P., Raschner, C., & Fink, C. (2019). Fitness, Knee Function and Competition Performance in Professional Alpine Skiers after ACL Injury. *Journal of Science and Medicine in Sport*, 22, S39–S43. <https://doi.org/10.1016/j.jsams.2018.06.014>
- Dluzniewski, A., Casanova, M. P., Ullrich-French, S., Brush, C. J., Larkins, L. W., & Baker, R. T. (2024). Psychological Readiness for Injury Recovery: Evaluating Psychometric Properties of the IPRRS and Assessing Group Differences in Injured Physically Active Individuals. *BMJ Open Sport and Exercise Medicine*, 10(2), 1-9. <https://doi.org/10.1136/bmjsem-2023-001869>
- Du, T., Shi, Y., Huang, H., Liang, W., & Miao, D. (2022). Current study on the influence of psychological factors on returning to sports after ACLR. In *Heliyon*, 8(12), 1-9. Elsevier Ltd. <https://doi.org/10.1016/j.heliyon.2022.e12434>
- Faleide, A. G. H., Magnussen, L. H., Bogen, B. E., Strand, T., Mo, I. F., Vervaat, W., & Inderhaug, E. (2021). Association Between Psychological Readiness and Knee Laxity and Their Predictive Value for Return to Sport in Patients With Anterior Cruciate Ligament Reconstruction. *American Journal of Sports Medicine*, 49(10), 2599–2606. <https://doi.org/10.1177/03635465211021831>
- Juggath, C., & Naidoo, R. (2024). The Influence of Psychological Readiness of Athletes when Returning to Sport after Injury. *South African Journal of Sports Medicine*, 36(1), 1–5. <https://doi.org/10.17159/2078-516x/2024/v36i1a16356>
- Kanakamedala, A. C., Anderson, A. F., & Irrgang, J. J. (2016). IKDC Subjective Knee Form and Marx Activity Rating Scale are suitable to evaluate all orthopaedic sports medicine knee conditions: a systematic review. *Journal of ISAKOS*, 1(1), 25–31. <https://doi.org/10.1136/jisakos-2015-000014>
- Kay, J., Liotta, E. S., Sugimoto, D., & Heyworth, B. E. (2023). Assessment of Return to Sport after ACL Reconstruction with Soft Tissue Autograft in Adolescent Athletes: Quadriceps Versus Hamstring Tendon. *Orthopaedic Journal of Sports Medicine*, 11(11), 1-8. <https://doi.org/10.1177/23259671231207113>

- Khan, T., Alvand, A., Prieto-Alhambra, D., Culliford, D. J., Judge, A., Jackson, W. F., Scammell, B. E., Arden, N. K., & Price, A. J. (2019). ACL and Meniscal Injuries Increase the Risk of Primary Total Knee Replacement for Osteoarthritis: A Matched Case-Control Study using the Clinical Practice Research Datalink (CPRD). *British Journal of Sports Medicine*, 53(15), 965–968. <https://doi.org/10.1136/bjsports-2017-097762>
- Lee, O. T., Williams, M. A., Shaw, C. D., & Delextrat, A. (2023). The Role of Strength-Related Factors on Psychological Readiness for Return to Sport Following Anterior Cruciate Ligament (ACL) Reconstruction. *Healthcare (Switzerland)*, 11(20), 1-12. <https://doi.org/10.3390/healthcare11202787>
- Lindanger, L., Strand, T., Mølster, A. O., Solheim, E., & Inderhaug, E. (2019). Return to Play and Long-term Participation in Pivoting Sports After Anterior Cruciate Ligament Reconstruction. *American Journal of Sports Medicine*, 47(14), 3339–3346. <https://doi.org/10.1177/0363546519878159>
- Meierbachtol, A., Yungtum, W., Paur, E., Bottoms, J., & Chmielewski, T. L. (2018). Psychological and Functional Readiness for Sport Following Advanced Group Training in Patients with Anterior Cruciate Ligament Reconstruction. *Journal of Orthopaedic and Sports Physical Therapy*, 48(11), 864–872. <https://doi.org/10.2519/jospt.2018.8041>
- Nawasreh, Z., Logerstedt, D., Cummer, K., Axe, M., Risberg, M. A., & Snyder-Mackler, L. (2018). Functional Performance 6 Months after ACL Reconstruction can Predict Return to Participation in the Same Preinjury Activity Level 12 and 24 Months after Surgery. *British Journal of Sports Medicine*, 52(6), 375. <https://doi.org/10.1136/bjsports-2016-097095>
- Nedder, V. J., Raju, A. G., Moyal, A. J., Calcei, J. G., & Voos, J. E. (2025). Impact of Psychological Factors on Rehabilitation After Anterior Cruciate Ligament Reconstruction: A Systematic Review. *Sports health*, 17(2), 291–298. <https://doi.org/10.1177/19417381241256930>
- Norte, G. E., Solaas, H., Saliba, S. A., Goetschius, J., Slater, L. V., & Hart, J. M. (2019). The Relationships between Kinesiophobia and Clinical Outcomes after ACL Reconstruction Differ by Self-Reported Physical Activity Engagement. *Physical therapy in sport : official journal of the Association of Chartered Physiotherapists in Sports Medicine*, 40, 1–9. <https://doi.org/10.1016/j.ptsp.2019.08.002>
- Nwachukwu, B. U., Adjei, J., Rauck, R. C., Chahla, J., Okoroha, K. R., Verma, N. N., Allen, A. A., & Williams, R. J. (2019). How Much Do Psychological Factors Affect Lack of Return to Play After Anterior Cruciate Ligament Reconstruction? A Systematic Review. *Orthopaedic journal of sports medicine*, 7(5), 2325967119845313. <https://doi.org/10.1177/2325967119845313>
- Pamukoff, D. N., Pietrosimone, B. G., Ryan, E. D., Lee, D. R., & Blackburn, J. T. (2017). Quadriceps Function and Hamstrings Co-Activation after Anterior Cruciate Ligament Reconstruction. *Journal of athletic training*, 52(5), 422–428. <https://doi.org/10.4085/1062-6050-52.3.05>
- Sadeqi, M., Klouche, S., Bohu, Y., Herman, S., Lefevre, N., & Gerometta, A. (2018). Progression of the Psychological ACL-RSI Score and Return to Sport After Anterior Cruciate Ligament Reconstruction: A Prospective 2-Year Follow-up Study From the French Prospective Anterior Cruciate Ligament Reconstruction Cohort Study (FAST). *Orthopaedic Journal of Sports Medicine*, 6(12), 1-7. <https://doi.org/10.1177/2325967118812819>
- Schilaty, N. D., McPherson, A. L., Nagai, T., & Bates, N. A. (2023). Differences in Psychological Readiness for Return to Sport after Anterior Cruciate Ligament Injury is Evident in Thigh Musculature Motor Unit Characteristics. *BMJ Open Sport and Exercise Medicine*, 9(3), 1-10. <https://doi.org/10.1136/bmjsem-2023-001609>

- Stigert, M., Ashnai, F., Thomeé, R., Hamrin Senorski, E., & Beischer, S. (2023). Physical Inactivity 5-8 Years after Anterior Cruciate Ligament Reconstruction is Associated with Knee-Related Self-Efficacy and Psychological Readiness to Return to Sport. *BMJ Open Sport and Exercise Medicine*, 9(4), 1-7. <https://doi.org/10.1136/bmjsem-2023-001687>
- Vutescu, E. S., Orman, S., Garcia-Lopez, E., Lau, J., Gage, A., & Cruz, A. I. (2021). Psychological and Social Components of Recovery Following Anterior Cruciate Ligament Reconstruction in Young Athletes: A Narrative Review. *International Journal of Environmental Research and Public Health*, 18(17), 1-8. <https://doi.org/10.3390/ijerph18179267>
- Webster, K. E., & Feller, J. A. (2019). Clinical Tests Can Be Used to Screen for Second Anterior Cruciate Ligament Injury in Younger Patients Who Return to Sport. *Orthopaedic Journal of Sports Medicine*, 7(8), 1-7. <https://doi.org/10.1177/2325967119863003>
- Webster, K. E., Nagelli, C. V., Hewett, T. E., & Feller, J. A. (2018). Factors Associated with Psychological Readiness to Return to Sport after Anterior Cruciate Ligament Reconstruction Surgery. *American Journal of Sports Medicine*, 46(7), 1545–1550. <https://doi.org/10.1177/0363546518773757>
- Yuliana, E., & Kushartanti, D. W. (2020). Knee Functional and Psychological Readiness of Post ACL Injury in Operative and Non-Operative Treatment. *Jurnal SPORTIF: Jurnal Penelitian Pembelajaran*, 6(3), 561–574. https://doi.org/10.29407/js_unpgri.v6i3.14626